

SAFETY DIRECTIONS

Product is poisonous if swallowed. Will irritate the eyes and skin. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhaled vapour or spray mist. When opening the container and preparing spray or wash, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length PVC gloves and water-resistant footwear. When using the prepared spray or wash, wear cotton overalls to the neck and wrist (or equivalent clothing), a washable hat, elbow-length PVC gloves and water-resistant footwear. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131 126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, do NOT induce vomiting. Give a glass of water. Activated charcoal may be advised. Give atropine if instructed.



9 338580 002000



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



Nucidol[®] 200 EC

Insecticide and Acaricide

Active Constituent: 200 g/L DIAZINON (AN ANTICHOLINESTERASE COMPOUND)
Solvent: 552 g/L HYDROCARBON-LIQUID

**Controls cattle lice and buffalo fly* on cattle, mange and lice on pigs,
flies and lice on horses, lice on goats, & flies in animal sheds.**

**ORGANOPHOSPHOROUS
PESTICIDES LIQUID, TOXIC,
N.O.S. UN NO. 3018
PACKAGING GROUP III**

*Diazinon sensitive buffalo flies



Contents: 500 mL

ZAGRO

READ ATTACHED LEAFLET AND THE DIRECTIONS FOR USE BEFORE OPENING
OR USING THIS PRODUCT

WITHHOLDING PERIODS:

MEAT (CATTLE AND HORSES): DO NOT USE less than 3 days before slaughter for human consumption.

MEAT (PIGS AND GOATS): DO NOT USE less than 14 days before slaughter for human consumption.
MILK (CATTLE AND GOATS): DO NOT USE on lactating or pregnant cows or nannies, where milk or milk products may be used for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact Pacific Biologics on 07 3283 5077 before using this product.

STORAGE

Store in original container tightly closed in a dry cool place. DO NOT store in direct sunlight. Store below 30°C (Room Temperature).

DISPOSAL

Triple or (preferably) pressure rinse containers into spray or other prepared solution. Do not dispose of undiluted chemicals on-site. Break, crush, or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned.

WARRANTY AND EXCLUSION OF LIABILITY

This product is warranted fit for the purposes specifically recommended by Zagro Animal Health Pte Ltd when used strictly as directed on this label. All other warranties and obligations or liabilities, whether expressed or implied by statute or otherwise, are excluded to the full extent that exclusion is permitted by law.

AUSTRALIAN DISTRIBUTOR:

Pacific Biologics Pty Ltd
35 Beach Street, Kippa-Ring, QLD 4021. ph: 07 3283 5077 or <http://www.pacificbiologics.com.au>

ZAGRO ANIMAL HEALTH PTE LTD
Zagro Global Hub, 5 Woodlands Terrace
Singapore 738430

ph: 0011 65 6759 1811 fax: 0011 65 6759 1855

APVMA Approval No. 49876/500mL/0509

Batch Number	
Expiry Date	

File name	49876_46891_500ml_Primary_MPL_V03 .pdf
Dimensions	(W) 27 cm x (H) 9.5 cm
Scale	100%
Date	26/02/10, V03

POISON

KEEP OUT OF REACH OF CHILDREN

Nucidol[®] 200 EC

INSECTICIDE AND ACARICIDE

Active Constituent:
200 g/L DIAZINON
(AN ANTICHOLINESTERASE COMPOUND)

Solvent:
552 g/L HYDROCARBON-LIQUID



Controls cattle lice and buffalo fly* on cattle, mange and lice on pigs, flies and lice on horses, lice on goats, & flies in animal sheds.

*Diazinon sensitive buffalo flies

READ THIS LEAFLET AND THE DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT

File name	49876_46891_500ml_Leaflet_MPL_V03.pdf
Dimensions	(W) 23.4 cm x (H) 33.2 cm
Scale	70%
Date	23/02/10, V03

DIRECTIONS FOR USE:

RESTRAINTS

DO NOT use on poultry.
DO NOT use on cats.
DO NOT use on lactating or pregnant cows or nannies, where milk or milk products may be used for human consumption.
DO NOT allow water to enter this container.
DO NOT rinse lid with water.

Animal/Situation	Pest	Rate		Withholding Period	Critical Comments
		High Volume	Other Application		
Cattle	Lice (<i>Bovicola bovis</i> ; <i>Linognathus vituli</i>)	250 mL per 100 litres of water	-	3 days	Apply at least 4.5 litres of spray per beast using hand spraying or high volume cattle spray units. Ensure animals is thoroughly covered. Respray as necessary.
		-	Low volume cattle sprayer: 500 mL per 100 litres of water		Apply at least 2.3 litres of spray per beast using low volume cattle sprayer. Examine cattle after 17 days for newly hatched lice and later during the season for further lice activity. Respray as necessary.
	Buffalo Fly (<i>Haematobia irritans exigua</i>)	400 mL per 100 litres of water	-		Apply 500 mL per beast and spray along backline. Respray as necessary.
		-	Backrubber: 500 mL per 10 litres of oil		Soak backrubber thoroughly. Repeat treatment in 14 days.
Pigs	Lice (<i>Haematopinus suis</i>)	250 mL per 100 litres of water	-	14 days	Spray pigs and sties thoroughly. Repeat treatment in 14 days.
					Mange (<i>Sarcoptes scabiei</i>)
Goats	Biting and Sucking Lice (<i>Damalina caprae</i> ; <i>Linognathus stenopsis</i>)	250 mL per 100 litres of water	-		Spray goats thoroughly making sure hair is wet to the skin. Examine goats after 14 days for newly hatched lice and later during the season for further lice activity. Respray as necessary.
Horses	Flies (<i>Musca vetustissima</i>) and Lice (<i>Damalina equi</i> ; <i>Haematopinus asini</i>)	25 mL per 10 litres of water	Swab: 25 mL per 10 litres of water	3 days	Spray or swab liberally as required.
Animal Sheds	Flies (<i>Musca domestica</i> ; <i>Stomoxys calcitrans</i>)	250 mL per 10 litres of water	-	-	Spray the inner walls thoroughly and any other places where flies settle. Respray as necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:
MEAT (CATTLE AND HORSES): DO NOT USE less than 3 days before slaughter for human consumption.
MEAT (PIGS AND GOATS): DO NOT USE less than 14 days before slaughter for human consumption.
MILK (CATTLE AND GOATS): DO NOT USE on lactating or pregnant cows or nannies, where milk or milk products may be used for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact Pacific Biologics on 07 3283 5077 before using this product.

GENERAL INSTRUCTIONS

NUCIDOL mixes readily with hard or soft water.

SAFETY DIRECTIONS

Product is poisonous if swallowed. Will irritate the eyes and skin. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When opening the container and preparing spray or wash, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length PVC gloves and water-resistant footwear. When using the prepared spray or wash, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length PVC gloves and water-resistant footwear. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Center (Phone Australia 131 126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, do NOT induce vomiting. Give a glass of water. Activated charcoal may be advised. Give atropine if instructed.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required, refer to Material Safety Data Sheet, which is available from <http://www.pacificbiologics.com.au>.

Protection of Wildlife, Fish, Crustacea and Environment

DO NOT contaminate streams, rivers or waterways with the chemical, waste liquid or containers.

STORAGE

Store in original container tightly closed in a dry cool place. DO NOT store in direct sunlight. Store below 30°C (Room Temperature).

DISPOSAL

Triple or (preferably) pressure rinse containers into spray or other prepared solution. Do not dispose of undiluted chemicals on-site. Break, crush, or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned.

WARRANTY AND EXCLUSION OF LIABILITY:

This product is warranted fit for the purposes specifically recommended by Zagro Animal Health Pte Ltd, Singapore when used strictly as directed in this leaflet. All other warranties and obligations or liabilities, whether expressed or implied by statute or otherwise, are excluded to the full extent that exclusion is permitted by law.

AUSTRALIAN DISTRIBUTOR:

Pacific BioLogics Pty Ltd
 35 Beach Street
 Kippa-Ring, QLD 4021
 ph (07) 3283 5077
www.pacificbiologics.com.au



ZAGRO ANIMAL HEALTH PTE LTD

Zagro Global Hub
 5 Woodlands Terrace
 Singapore 738430
 ph: 65 6759 1811 fax: 65 6759 1855
zsingapore@zagro.com
www.zagro.com

APVMA Approval Number: 49876/500 mL/0509

® Registered trademark of Zagro Singapore Pte Ltd

File name	49876_46891_500ml_Leaflet_MPL_V03.pdf
Dimensions	(W) 23.4 cm x (H) 33.2 cm
Scale	70%
Date	23/02/10, V03